

# NEXT-LEVEL PARASITE PROTECTION



FLEAS



TICKS



HEARTWORM DISEASE



ROUNDWORMS



HOOKWORMS



## OPTIMIZED MOXIDECTIN DOSE >>

### DESIGNED WITH EFFICACY AND SAFETY IN MIND

#### Minimum recommended dose of 12 mcg/kg monthly

- This dose is higher than the dose historically shown to be effective against susceptible isolates
- Provides a wide margin of safety for MDR1-deficient dogs
- In the Avermectin-Sensitive Collie Safety study, no clinical signs of avermectin toxicity were noted in any dog at any time

#### Recommended monthly dosing for at least 6 months after last exposure to mosquitoes

- All NexGard® PLUS (afoxolaner, moxidectin, and pyrantel chewable tablets) pivotal heartworm disease studies were conducted using 6 monthly doses after infection or last exposure to mosquitoes
- NexGard PLUS chews were 100% effective in all heartworm disease studies that were performed for approval of the product (2 laboratory studies and 1 field study)
- This recommendation more closely aligns with the standard recommendation for heartworm disease preventives to be given year-round and is more representative of how the products are used in the real world
- **While the dose is an important factor, administering sequential, oral monthly doses profoundly increases the effectiveness against rare, less-susceptible heartworm isolates.**

**IMPORTANT SAFETY INFORMATION:** NexGard® PLUS chews are for use in dogs only. The most frequently reported adverse reactions reported in clinical trials were diarrhea, vomiting, lethargy, and itching. NexGard PLUS contains afoxolaner, a member of the isoxazoline class. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures in dogs with or without a history of seizures. Use with caution in dogs with a history of seizures or neurologic disorders. The safe use of NexGard PLUS has not been evaluated in breeding, pregnant, or lactating dogs. Dogs should be tested for existing heartworm infection prior to starting a heartworm disease preventive. For more information, see full prescribing information or visit [NexGardPLUSClinic.com](http://NexGardPLUSClinic.com).



NexGard® is a registered trademark of Boehringer Ingelheim Animal Health France, used under license. All other trademarks are the property of their respective owner. ©2025 Boehringer Ingelheim Animal Health USA Inc., Duluth, GA. All rights reserved. US-PET-0594-2023-V4.

**NexGard® PLUS**  
(afoxolaner, moxidectin, and  
pyrantel chewable tablets)

# AN OPTIMIZED DOSE THAT IS EFFECTIVE AND SAFE ➤

## Oral moxidectin in dogs<sup>1</sup>:

- First moxidectin-containing oral heartworm disease preventive was approved in 1997
- Recommended dose of 3 mcg/kg monthly
- In lab studies, oral moxidectin dosed as low as 0.5 mcg/kg was 100% effective

## Identification of heartworm resistance<sup>2-4</sup>:

- In the past 2 decades, several less-susceptible *Dirofilaria immitis* isolates have been identified
- Resistant subpopulations of heartworms appear to be an infrequent, focal issue seen in localized regions of the country and do not appear to be widespread


## Effectiveness against rare, less-susceptible isolates can be improved by:

- Increasing the moxidectin mcg/kg dose
- Increasing the number of monthly oral administrations

## Efficacy of a single administration of moxidectin at different doses against the JYD-34 heartworm isolate

Increasing the moxidectin mcg/kg dose improved activity against this less-susceptible heartworm isolate. However, the effectiveness of a single 24 mcg/kg dose was still poor.

Even with a single dose as high as 100 mcg/kg, 100% efficacy was not achieved. Additionally, this high of a moxidectin dose would not allow for an adequate margin of safety in MDR1-deficient dogs.



- █ Data from Zoetis published studies\*   █ Data from Boehringer Ingelheim internal studies†
- 3 mcg/kg = historically 100% effective against susceptible isolates
  - 12 mcg/kg = minimum recommended dose in *NexGard® PLUS* (afoxolaner, moxidectin, and pyrantel chewable tablets)
  - 24 mcg/kg = minimum recommended dose in *Simparica Trio®* (sarolaner, moxidectin, and pyrantel chewable tablets)

\* Savadelis MD, McTier TL, Kryda K, Maeder SJ, Woods DJ. Moxidectin: heartworm disease prevention in dogs in the face of emerging macrocyclic lactone resistance. *Parasit Vectors*. 2022;15:82.

† Martin EM, Mitchell EB, Yoon S, McCall JW, et al. Efficacy of moxidectin, using various dose regimens, against JYD-34, a macrocyclic lactone resistant isolate of *Dirofilaria immitis*. *Parasit Vectors*. 2024 Apr 4;17(1):176.

**IMPORTANT SAFETY INFORMATION:** *NexGard® PLUS* (afoxolaner, moxidectin, and pyrantel chewable tablets) is for use in dogs only. The most frequently reported adverse reactions reported in clinical trials were diarrhea, vomiting, lethargy, and itching. *NexGard PLUS* contains afoxolaner, a member of the isoxazoline class. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures in dogs with or without a history of seizures.


# NexGard® PLUS

(afoxolaner, moxidectin, and pyrantel chewable tablets)



**While increasing the oral moxidectin mcg/kg dose improved effectiveness, increasing the number of monthly oral administrations profoundly increased effectiveness.**

## Efficacy of multiple monthly oral administrations of moxidectin given at different doses against the JYD-34 heartworm isolate across different studies



Differences above (solid bar, bar with lines and bar with dots) indicate different studies conducted at different times.

The dose and the number of doses administered vary in these data.

- 3 monthly doses at 3 mcg/kg were 44.4% effective
- 5 monthly doses at 6 mcg/kg were 83.3% and 90.2% effective
- Multiple monthly doses at 9 – 30 mcg/kg were ≥ 94.1% effective

This information is summarized from various studies over time, conducted at different laboratories. The moxidectin formulations used in these studies varied as well (oral solutions vs. shaved tablets). These data cannot be directly compared, because these are not head-to-head studies, and several factors unrelated to treatment can impact the results.

NexGard® PLUS chews were not used in any of the highlighted studies.

<sup>1</sup>Savadelis MD, McTier TL, Kryda K, Maeder SJ, Woods DJ. Moxidectin: heartworm disease prevention in dogs in the face of emerging macrocyclic lactone resistance. *Parasit Vectors*. 2022;15:82.

<sup>2</sup>Martin EM, Mitchell EB, Yoon S, McCall JW, et al. Efficacy of moxidectin, using various dose regimens, against JYD-34, a macrocyclic lactone resistant isolate of *Dirofilaria immitis*. *Parasit Vectors*. 2024 Apr 4;17(1):176.

<sup>3</sup>Freedom of Information Summary, PROHEART® Tablets, NADA 141-051, May 1997.

<sup>4</sup>Bourguinat C, Lee A, Lizundia R, et al. ML resistance in *D. immitis*: failure of heartworm preventives and investigation of genetic markers for resistance. *Vet Parasit*. 2015;210(3-4):167-178.

<sup>5</sup>American Heartworm Society Resistance Statement. American Heartworm Society. [https://d3ft8scckhnqjm2.cloudfront.net/images/Resistance\\_Statement\\_20200716.pdf](https://d3ft8scckhnqjm2.cloudfront.net/images/Resistance_Statement_20200716.pdf). Accessed May 1, 2023.

<sup>6</sup>Pulaski C, Malone J, Bourguinat C, et al. Establishment of macrocyclic lactone resistant *Dirofilaria immitis* isolates in experimentally infected laboratory dogs. *Parasit Vectors*. 2014;7:7494.

Use with caution in dogs with a history of seizures or neurologic disorders. The safe use of *NexGard PLUS* has not been evaluated in breeding, pregnant, or lactating dogs. Dogs should be tested for existing heartworm infection prior to starting a heartworm disease preventive. For more information, see full prescribing information or visit [NexGardPLUSClinic.com](http://NexGardPLUSClinic.com).

# NexGard® PLUS

## (afoxolaner, moxidectin, and pyrantel chewable tablets)

### For oral use in dogs only.

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

#### Description:

NexGard® PLUS (afoxolaner, moxidectin, and pyrantel chewable tablets) is available in five sizes of beef-flavored, soft chewables for oral administration to dogs and puppies according to their weight. Each chewable is formulated to provide minimum doses of 1.14 mg/lb (2.5 mg/kg) afoxolaner, 5.45 mcg/lb (12 mcg/kg) moxidectin, and 2.27 mg/lb (5.0 mg/kg) pyrantel (as pamoate salt).

Afoxolaner is a member of the isoxazoline family of compounds. Its chemical name is 1-Naphthalene-carboxamide,4-[5-[3-chloro-5-(trifluoromethyl)-phenyl]-4,5-dihydro-5-(trifluoromethyl)-3-isoxazolyl]-N-[2-oxo-2-[(2,2,2-trifluoroethyl)amino]ethyl].

Moxidectin is a semisynthetic macrocyclic lactone derived from the actinomycete *Streptomyces cyanescens* noncyrogenus. The chemical name for moxidectin is [6R,23E,25S](E)-5-O-Demethyl-28-deoxy-25-(1,3-dimethyl-1-butene)-6,28-epoxy-23-(methoxyimino) milbemycin B.

Pyrantel is a member of the tetrahydropyrimidine family of compounds. Its chemical name is (E)-1,4,5,6-Tetrahydro-1-methyl-2-[2-(2-thienyl) vinyl] pyrimidine 4, 4' methylenebis [3-hydroxy-2-naphthoate](1:1).

#### Indications:

NexGard® PLUS is indicated for the prevention of heartworm disease caused by *Dirofilaria immitis* and for the treatment and control of adult hookworm (*Ancylostoma caninum*, *Ancylostoma braziliense*, and *Uncinaria stenocephala*) and roundworm (*Toxocara canis* and *Toxascaris leonina*) infections.

NexGard® PLUS kills adult fleas and is indicated for the treatment and prevention of flea infestations (*Ctenocephalides felis*) and the treatment and control of *Ixodes scapularis* (black-legged tick), *Rhipicephalus sanguineus* (brown dog tick), *Dermacentor variabilis* (American dog tick), *Amblyomma americanum* (lone star tick), and *Haemaphysalis longicornis* (longhorned tick) infestations for one month in dogs and puppies eight weeks of age and older, weighing four pounds of body weight or greater.

#### Dosage and Administration:

NexGard® PLUS is given orally once a month at the minimum dosage of 1.14 mg/lb (2.5 mg/kg) afoxolaner, 5.45 mcg/lb (12 mcg/kg) moxidectin, and 2.27 mg/lb (5.0 mg/kg) pyrantel (as pamoate salt).

**For heartworm disease prevention, give once monthly for at least six months after last exposure to mosquitoes (see Effectiveness).**

#### Dosing Schedule:

Body Weight (lbs.)	Afoxolaner Per Chewable (mg)	Moxidectin Per Chewable (mcg)	Pyrantel* Per Chewable (mg)	Chewables Administered
4 – 8	9.375	45	18.75	One
8.1 – 17	18.75	90	37.5	One
17.1 – 33	37.5	180	75	One
33.1 – 66	75	360	150	One
66.1 – 132	150	720	300	One
Over 132	Administrator	the appropriate combination of chewables		

\*As pamoate salt.

NexGard® PLUS can be administered with or without food. Care should be taken to ensure that the dog consumes the complete dose and that part of the dose is not lost or refused. If a dose is missed, administer NexGard® PLUS and resume a monthly dosing schedule.

#### Heartworm Prevention:

NexGard® PLUS should be administered at monthly intervals year-round or, at a minimum, administration should start within one month of the dog's first seasonal exposure to mosquitoes and should continue at monthly intervals until at least six months after the dog's last exposure (see Effectiveness). When replacing another monthly heartworm preventive product, the first dose of NexGard® PLUS should be given within a month of the last dose of the former medication.

#### Flea Treatment and Prevention:

NexGard® PLUS should be administered year-round at monthly intervals or started at least one month before fleas become active. To minimize the likelihood of flea reinfection, it is important to treat all animals within a household with an approved flea control product.

#### Tick Treatment and Control:

NexGard® PLUS should be administered year-round at monthly intervals or started at least one month before ticks become active.

#### Intestinal Nematode Treatment and Control:

NexGard® PLUS treats and controls adult hookworms (*Ancylostoma caninum*, *Ancylostoma braziliense*, and *Uncinaria stenocephala*) and roundworms (*Toxocara canis* and *Toxascaris leonina*). For the treatment of adult hookworm and roundworm infections, NexGard® PLUS should be administered as a single dose. Monthly use of NexGard® PLUS will control any subsequent infections. Dogs may be exposed to and can become infected with hookworms and roundworms throughout the year, regardless of season or climate.

#### Contraindications:

There are no known contraindications for the use of NexGard® PLUS.

#### Warnings:

Not for use in humans. Keep this and all drugs out of the reach of children. In case of accidental ingestion, contact a physician for treatment advice.

Keep NexGard® PLUS in a secure location out of the reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

#### Precautions:

Afoxolaner, one of the ingredients in NexGard® PLUS, is a member of the isoxazoline class. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures. Seizures have been

reported in dogs receiving isoxazoline class drugs, even in dogs without a history of seizures. Use with caution in dogs with a history of seizures or neurologic disorders.

Treatment with fewer than six monthly doses after the last exposure to mosquitoes has not been evaluated and may not provide complete heartworm prevention.

Prior to administration of NexGard® PLUS, dogs should be tested for existing heartworm infection. At the discretion of the veterinarian, infected dogs should be treated with an adulticide to remove adult heartworms. NexGard® PLUS is not effective against adult *D. immitis*.

The safe use of NexGard® PLUS in breeding, pregnant, or lactating dogs has not been evaluated.

#### Adverse Reactions:

In a field safety and effectiveness study, NexGard® PLUS was administered to dogs for the prevention of heartworm disease. The study included a total of 272 dogs (134 administered NexGard® PLUS and 138 administered active control) treated once monthly for 11 treatments.

Over the 330-day study period, all observations of potential adverse reactions were recorded. The most frequent reactions reported in the NexGard® PLUS group are presented in the following table.

**Table 1: Dogs With Adverse Reactions.**

Clinical Sign	NexGard® PLUS n = 134 Number (Percentage)	Active Control n = 138 Number (Percentage)
Diarrhea	9 (6.7%)	7 (5.1%)
Vomiting	6 (4.5%)	7 (5.1%)
Lethargy	3 (2.2%)	5 (3.6%)
Itching	3 (2.2%)	3 (2.2%)
Dermatitis	2 (1.5%)	1 (0.7%)
Anorexia	1 (0.7%)	4 (2.9%)
Muscle tremor	1 (0.7%)	1 (0.7%)

One dog in the NexGard® PLUS group was reported to exhibit muscle tremors along with nausea and depression for one day after the Day 0 treatment. The dog remained in the study and muscle tremors were not reported after any subsequent treatments.

#### Contact Information:

For a copy of the Safety Data Sheet (SDS) or to report suspected adverse drug events, contact Boehringer Ingelheim Animal Health USA Inc. at 1-888-637-4251 or [www.nexgardforpets.com](http://www.nexgardforpets.com).

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or [www.fda.gov/reportanadverse](http://www.fda.gov/reportanadverse).

#### Clinical Pharmacology:

##### Mode of Action:

NexGard® PLUS (afoxolaner, moxidectin, and pyrantel chewable tablets) contains the three active pharmaceutical ingredients afoxolaner, moxidectin, and pyrantel (as pamoate salt).

Afoxolaner is a member of the isoxazoline family, shown to bind at a binding site to inhibit insect and acarine ligand-gated chloride channels, in particular those gated by the neurotransmitter gamma-aminobutyric acid (GABA), thereby blocking pre- and postsynaptic transfer of chloride ions across cell membranes. Prolonged afoxolaner-induced hyperexcitation results in uncontrolled activity of the central nervous system and death of insects and acarines. The selective toxicity of afoxolaner between insects and acarines and mammals may be inferred by the differential sensitivity of the insects and acarines GABA receptors versus mammalian GABA receptors.

Moxidectin is an endectocide in the macrocyclic lactone class. Moxidectin acts by interfering with chloride channel-mediated neurotransmission in susceptible parasites, which results in paralysis and death of the parasite.

Pyrantel is a nematocide belonging to the tetrahydropyrimidine class. Pyrantel acts as a depolarizing, neuromuscular-blocking agent in susceptible parasites, causing paralysis and death or expulsion of the parasite.

#### Pharmacokinetics:

Following a single oral administration of a near-final formulation of NexGard® PLUS (at mean doses of 3.9 mg/kg afoxolaner, 18.8 mcg/kg moxidectin, and 7.8 mg/kg pyrantel pamoate) in fed and fasted Beagle dogs (10 to 21 months of age), afoxolaner and moxidectin were more rapidly absorbed in the fasted state with a time to maximum concentration (t<sub>max</sub>) of 2 to 3 hours.

The afoxolaner mean maximum plasma concentrations (C<sub>max</sub>) in the fed and fasted states were 1610 and 2200 ng/mL (CV=33 and 16%) and the moxidectin mean C<sub>max</sub> values were 11.1 and 15.5 ng/mL (CV=39 and 24%), respectively. The area under the curve (AUC) for afoxolaner and moxidectin were similar between fed and fasted states. Post-dose pyrantel plasma concentrations were quantifiable out to 24 hours.

Following six oral administrations of NexGard® PLUS at 1, 3, and 5X the maximum exposure dose of 5 mg/kg, 24 mcg/kg, and 10 mg/kg afoxolaner, moxidectin, and pyrantel pamoate, respectively, every 28 days in 8-week-old Beagle dogs, afoxolaner and moxidectin t<sub>max</sub> ranged from 2 to 6 hours. The observed mean C<sub>max</sub> and AUC at steady state in the 1X dose group were 2230 ng/mL and 19000 days·ng/mL for afoxolaner and 14.8 ng/mL and 55.2 days·ng/mL for moxidectin, respectively. Based on mean C<sub>max</sub>, afoxolaner and moxidectin accumulated by less than 4-fold at steady state. Afoxolaner and moxidectin exposure increased in a dose proportional manner between the 1X and 3X dose groups but was less than dose proportional in the 5X dose group.

Pyrantel pamoate is poorly absorbed into systemic circulation. Pyrantel pamoate is intended to remain in the gastrointestinal tract to allow effective concentrations to be delivered to gastrointestinal nematodes.

#### Effectiveness:

##### Heartworm Prevention:

In two well-controlled laboratory studies, NexGard® PLUS was 100% effective against induced *D. immitis* infections when administered for six consecutive months.

In a well-controlled US field study consisting of 120 dogs administered NexGard® PLUS and 124 administered an active control, no dogs treated with NexGard® PLUS tested positive for heartworm disease. All dogs treated with NexGard® PLUS were negative for *D. immitis* antigen and blood microfilariae at study completion on Day 330.

#### Flea Treatment and Prevention:

In a well-controlled laboratory study, NexGard® PLUS demonstrated ≥99.8% effectiveness against adult fleas 24 hours after weekly infestations for one month.

In a separate well-controlled laboratory study, afoxolaner alone began to kill fleas four hours after initial administration and demonstrated >99% effectiveness at eight hours.

In an additional well-controlled laboratory study, afoxolaner alone demonstrated 100% effectiveness against adult fleas 24 hours post-infestation for 35 days and was ≥93% effective at 12 hours post-infestation through Day 21 and on Day 35. On Day 28, afoxolaner alone was 81.1% effective 12 hours post-infestation. Dogs in both the afoxolaner-treated and control groups that were infested with fleas on Day 1 generated flea eggs at 12 and 24 hours post-treatment (0-11 eggs and 1-17 eggs in the afoxolaner-treated dogs, and 4-90 eggs and 0-118 eggs in the control dogs, at 12 and 24 hours, respectively). At subsequent evaluations post-infestation, fleas from dogs in the afoxolaner-treated group were essentially unable to produce any eggs (0-1 eggs), while fleas from dogs in the control group continued to produce eggs (1-141 eggs).

In a 90-day US field study conducted in households with existing flea infestations of varying severity, the effectiveness of afoxolaner alone against fleas on the Day 30, 60, and 90 visits compared with baseline was 98.0%, 99.7%, and 99.9%, respectively. In a second 90-day US field study, the effectiveness of afoxolaner alone against fleas on the Day 30, 60, and 90 visits compared with baseline was 97.5%, 99.7%, and 99.9%, respectively. Dogs in the second study with signs of Flea Allergy Dermatitis (FAD) showed improvement in erythema, alopecia, papules, scales, crusts, and excoriation following treatment, as a direct result of eliminating fleas.

Collectively, the data from the five studies (three laboratory and two field) demonstrate that NexGard® PLUS kills fleas before they can lay eggs, thus preventing subsequent flea infestations after the start of treatment of existing flea infestations.

#### Tick Treatment and Control:

In well-controlled laboratory studies, NexGard® PLUS demonstrated ≥97% effectiveness against *Amblyomma americanum* 72 hours post-infestation, for one month.

In well-controlled laboratory studies, a chewable containing afoxolaner alone, one of the active ingredients in NexGard® PLUS, demonstrated effectiveness against *Ixodes scapularis*, *Rhipicephalus sanguineus*, and *Dermacentor variabilis* at 48 hours post-infestation, and against *Haemaphysalis longicornis* at 72 hours post-infestation, for one month.

#### Intestinal Nematode Treatment and Control:

Elimination of adult roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Ancylostoma braziliense*, and *Uncinaria stenocephala*) was demonstrated in well-controlled laboratory studies.

#### Target Animal Safety:

##### Margin of Safety:

NexGard® PLUS was administered orally at 1, 3, and 5X the maximum exposure doses at approximately 28-day intervals for six treatments to 8-week-old Beagle puppies. Dogs in the control group were sham-dosed. There were no clinically relevant, treatment-related effects on body weights, food consumption, clinical pathology (hematology, coagulation, serum chemistry, and urinalysis), gross pathology, histopathology, organ weights, or ophthalmic examinations. Mild, self-limiting diarrhea (with and without blood) was possibly related to treatment, as there were more incidences in the NexGard® PLUS groups than the control group throughout the study, including within 48 hours after treatment.

##### Avermectin-Sensitive Collie Safety:

NexGard® PLUS was administered orally at 1, 3, and 5X the maximum label dose to MDRI-deficient Collies once on Day 0, with a second administration to the 1X group on Day 28. Dogs in the control group were sham-dosed on Days 0 and 28. No clinical signs of avermectin toxicity were noted in any dog at any time during the study. Vomiting was observed in some dogs in the 3X and 5X groups and resolved without treatment. Diarrhea, with or without blood, was observed in some dogs in all of the NexGard® PLUS groups and resolved without treatment.

##### Heartworm-Positive Safety:

NexGard® PLUS was administered orally at 1X and 3X the maximum exposure doses at approximately 28-day intervals for three treatments to Beagle dogs with adult heartworm infections and circulating microfilariae. Dogs in the control group were sham-dosed. Diarrhea was observed in one dog in the 1X group and in three dogs in the 3X group, and vomiting was observed in two dogs in the 3X group. No signs of avermectin toxicity were observed at any time during the study. There were no clinical signs associated with death of the microfilariae observed in any of the dogs.

#### Field Safety:

In a well-controlled field study, NexGard® PLUS was used concurrently with other medications such as vaccines, antibiotics, non-steroidal anti-inflammatory drugs (NSAIDs), anesthetics, sedatives, analgesics, steroids, antihelmintics, antiemetics, and antipruritics. No adverse reactions were associated with the concurrent use of NexGard® PLUS and other medications.

#### How Supplied:

NexGard® PLUS is available in five strengths of beef-flavored soft chewables formulated according to the weight of the dog (see Dosage and Administration). Each chewable size is available in color-coded packages of 1, 3, or 6 chewables.

#### Storage Information:

Store in original package at or below 25°C (77°F) with excursions permitted up to 40°C (104°F).

Approved by FDA under NADA # 141-554

Marketed by: Boehringer Ingelheim Animal Health USA Inc., Duluth, GA 30096

NexGard® is a registered trademark of Boehringer Ingelheim Animal Health France, used under license.

©2024 Boehringer Ingelheim Animal Health USA Inc. All rights reserved. US-PET-0084-2023-V4.

181531-004  
Rev. 08/2024

